

GC2030

Golf Course 2030

The Climate Impact of Golf Courses

Carbon Balances in Golf Course
Landscapes (Version 2, 2025)



This report is part of the series of Golf Course 2030 publications on the climate impact of golf courses. The other reports in this series are:

- The Climate Impact of Golf Courses: A Case Study from Minchinhampton Golf Club, UK
- The Climate Impact of Golf Courses: A Sampling Protocol to Determine Golf Course Soil Carbon Fluxes

When estimating carbon sequestration, carbon emissions and the resulting carbon balance of golf courses it is critical to clearly define which parts of the golf facility are included in these calculations. Carbon emission estimates from golf courses could

include emissions from the clubhouse, event buildings and even from the transport of golfers and employees to the facility. Conversely, carbon sequestration estimates could include sequestration from non-turf areas of the golf course. In this series of reports, we focus solely on carbon emissions from golf course maintenance activities and carbon sequestration from golf course turfgrass surfaces.

Project conducted by:

Michael Bekken, PhD
Research Scientist
Norwegian Institute of Bioeconomy
michael.bekken@nibio.no

An introduction to carbon and climate	4	Practical steps	18
The basics of carbon emission and sequestration	6	Minimising carbon emissions	19
		Electrification of golf course maintenance equipment	20
Golf course carbon cycling	8	Maximising carbon sequestration	21
Carbon emissions in golf course turfgrass maintenance	9		
Carbon sequestration of turf and non-turf areas on golf courses	12	Summary	22
The carbon balance of golf courses	13	References	23
Golf course carbon cycle diagram	16		



An introduction to carbon and climate.

Since the industrial revolution, human activity has caused the average surface temperature of Earth to rise by approximately 1°C.

Anthropogenic emissions of greenhouse gases, primarily carbon dioxide, methane, nitrous oxide and fluorinated gases, which decrease the amount of heat Earth loses to space, are responsible for rising global average temperatures.

Throughout Earth's history, the climate has constantly, although slowly, changed. What is unprecedented today is the rate at which Earth's climate is changing. Human emissions of greenhouse gases are warming the climate so quickly, that life on earth is struggling to adapt to the rapidly changing conditions.

Global biodiversity is quickly declining, and we are under pressure to adapt to warmer temperatures, droughts, floods and increasingly powerful and destructive storms. The golf industry is not immune to these pressures and is experiencing a range of challenges and threats, from water shortages to heavy rainfall and floods. It is critical for the golf industry to contribute to mitigating climate change by reducing carbon emissions and maximising the sequestration value of golf courses.

It is critical for the golf industry to contribute to mitigating climate change by reducing carbon emissions and maximising carbon sequestration of courses.

This report uses “carbon emissions” as a shorthand for the production of greenhouse gases (GHGs). Most greenhouse gases contain carbon (eg carbon dioxide, methane, fluorinated gases) but not all do (eg nitrous oxide).



The basics of carbon emission and sequestration



Carbon emissions are the processes by which carbon is released into the atmosphere. Combustion of fossil fuels is the most common process that emits GHGs to the atmosphere and involves reacting oxygen and hydrocarbons (ie petrol, diesel, natural gas etc) to produce carbon dioxide and water.



Conversely, carbon sequestration is the process by which carbon is taken out of the atmosphere. Photosynthesis by plants removes carbon dioxide from the atmosphere and through a complex series of chemical reactions produces sugars and oxygen. The plant retains the sugars and releases the oxygen back into the atmosphere.

The plant-soil system is responsible for carbon sequestration in vegetated landscapes (ie forests, grasslands and golf courses etc). The sugars produced in photosynthesis are used by plants as food and incorporated into their tissues (ie wood, leaves etc). When plants die some of this plant material is consumed by microorganisms and other animals and the carbon in the plant tissue is returned to the atmosphere as CO₂. However, microorganisms cannot break down all plant tissue immediately, meaning that some of the plant tissue will slowly be transformed into what is called soil organic matter. Soil organic matter contains approximately 50-60% carbon

and is one of the main repositories of carbon on Earth. Carbon sequestration by plants largely happens through the combined processes of photosynthesis, eventual plant death and the decomposition of plant material into soil organic matter.

Soil organic matter, which stores carbon in soils, is not a permanent repository or reservoir of carbon. Soil microorganisms can eventually break down soil organic matter under certain conditions and, in doing so, return the carbon stored in soil organic matter into the atmosphere as CO₂.



In much the same way as we eat food that contains carbon and breathe out some of that carbon into the atmosphere as CO₂, soil microorganisms eat soil organic matter and release CO₂ back into the atmosphere. Scientists refer to this process performed by humans and microorganisms alike as respiration. Though the equation above is greatly simplified, respiration is essentially the opposite of photosynthesis.

Carbon, therefore, flows from the atmosphere into the soil and from the soil back into the atmosphere. The balance of these flows, which are always happening simultaneously, determines whether carbon accumulates in the soil or in the

atmosphere. Soils can be a great repository of carbon, but they can also be large sources of carbon to the atmosphere if they are disturbed, managed poorly or the climate changes.



Golf course carbon cycling.

Golf courses emit carbon into the atmosphere and sequester carbon from the atmosphere. The balance of emissions and sequestration determines the climate impact of a golf course.

The sum of direct, indirect and associated emissions are equal to the total carbon emissions from a golf course.

Carbon emissions in golf course turfgrass maintenance

Carbon emissions can be broken down into categories called 'scopes' based on how closely associated the emissions are with an organisation or activity. Most carbon accounting strategies use a three scope system to categorise emissions, which is described below in the context of golf course maintenance.

Scope 1 Direct emissions

Golf course turfgrass maintenance operations emit carbon directly into the atmosphere through activities such as the combustion of fuel to power mowers and other maintenance equipment. These emissions are referred to as direct or Scope 1 emissions because they are directly controlled by the golf course.

Scope 2 Indirect emissions

Golf courses also emit carbon indirectly through the use of electricity, which are referred to as Scope 2 emissions. Emissions from using electricity are not generated on the golf course but are emitted at a power plant where the electricity is generated, which is why these emissions are labelled as indirect.

The amount of carbon emitted per unit of electricity generated varies depending on the power source being used. Coal fired power plants, for example, generate more carbon emissions per unit of electricity than hydroelectric power plants.

Scope 3 Associated emissions

Golf courses not only emit carbon directly through burning fuels and indirectly via consuming electricity, but also through associated economic and supply chain related activities. The equipment (eg mowers, utility vehicles, tractors etc) and materials (eg fertiliser, pesticide, sand etc) that golf courses use all emit carbon in their production and transport to the facility. Emissions from these associated activities are referred to as Scope 3 emissions.

The sum of direct (Scope 1), indirect (Scope 2) and associated (Scope 3) emissions are equal to the total carbon emissions from a golf course.

Emissions category	Scope 1	Scope 2	Scope 3
Electricity 		Carbon emissions from the generation of grid electricity in the UK that is used at MGC.	Carbon emissions of electricity that is lost during the transmission and distribution of electricity in the UK.
Fertiliser 	Emissions from denitrification of nitrogen fertilisers after application.		Carbon emissions from the manufacturing of nitrogen, phosphorus and potassium fertilisers.
Fuel 	Carbon emissions from the combustion of all fuels at MGC. This includes fuels to power all machinery and heat the maintenance facility.		Carbon emissions from the manufacturing of fuels used at MGC.
Machinery 			Carbon emissions from the manufacturing, transport and repair of machinery at MGC.
Pesticide 			Carbon emissions from the manufacturing of herbicides, insecticides and fungicides used at MGC.
Sand 			Carbon emissions from the mining and transport of sand to the facility.

Study of golf course maintenance emissions in US and Europe (Part 1)

Recent research conducted by turfgrass scientists at University of Wisconsin-Madison, found that average carbon emissions on four US golf courses were approximately 4000kg CO₂e ha⁻¹ yr⁻¹ (Bekken and Soldat, 2021). Scope 1, 2 and 3 emissions were all similar, approximately 1300kg CO₂e ha⁻¹ yr⁻¹ each (Figure 1A). On three European golf courses, overall emissions levels were similar, approximately 4300kg CO₂e ha⁻¹ yr⁻¹ (Figure 1B). Scope 1 emissions on European golf courses were 1200kg CO₂e ha⁻¹ yr⁻¹, meaning they were approximately the same as in the US. Scope 2 emissions were much lower in Europe than in the US because of lower emissions from electricity

generation in Europe. However, Scope 3 emissions on the European golf courses in the study were much higher because of sand topdressing use on fairways at the three golf courses in Europe that participated in the study. The authors of the study recognise that topdressing fairways is not a standard practice in golf course maintenance in Europe. The results of the study highlight that if fairways are topdressed, the carbon cost to produce, transport and apply that sand will be high.

Figures 1C and D show the same four US golf courses and three European golf courses, but instead of categorising the emissions by scope, the emissions are categorised into Electricity (production and transport), Electricity (use), Fertiliser (production and

transport), Fuel (use), Machinery (production, transport, repair), Pesticide (production), and Sand (production and transport). For the four US golf courses Electricity (use) and Fuel (use) accounted for 63% of carbon emissions. For the three European golf courses, emissions were highest from Sand (production and transport) and Fuel (use).

The study revealed that if a golf course is not topdressing fairways, the most effective way to reduce carbon emissions is to transition to electric maintenance equipment and to source low carbon electricity.






Figure 1. Carbon emissions by scope on: A) four US golf courses and B) three European golf courses. Carbon emissions across eight emission categories on: C) four US golf courses and D) three European golf courses. P-Production, U-Use, A-Application (ie, fertiliser denitrification), T-Transport, and R-Repair.

The carbon balance of a golf course is equal to the weight of carbon emitted minus the weight of carbon sequestered.

Carbon sequestration of turf and non-turf areas on golf courses

Turfgrass sequestration

The turf-soil system on golf courses sequesters carbon in much the same way as natural vegetated landscapes. Golf courses are commonly converted from agricultural land and most forms of modern agriculture cause a decline in soil organic matter. When a golf course is constructed on formerly agricultural land, several scientific studies indicate that the turfgrass-soil system quickly sequesters carbon (ie soil organic matter levels go up) for the first 20-30 years after the golf course is built. However, after 30 years, the rate of carbon sequestration (ie the rise in soil organic matter) slows (Qian and Follet, 2002; Selhorst and Lal, 2011). After approximately 75 years, the rate of carbon sequestration becomes very low as soil organic matter levels stabilise and come to equilibrium. At this point, most studies suggest that the turf-soil system on golf courses still sequesters carbon, but at a low rate.

Sequestration in non-turf areas

Approximately 20% of the average US golf course is natural or native vegetation (ie, grassland, forest or wetland) (GCSAA, 2017). Areas of native vegetation are important components of a golf course for the biodiversity and habitat value that they provide, especially in urban settings.

These vegetated native areas of the golf course can also sequester carbon. However, the rate of carbon sequestration in native areas depends on the type of vegetation and how or if the native area is managed. The few scientific studies that attempt to account for carbon sequestration in native areas are mostly concerned in quantifying the carbon sequestration by forested areas on the golf course. The rate of forest carbon sequestration is heavily dependent on forest age and health, therefore the estimates of carbon sequestration in forested areas on golf courses must be carefully conducted if the estimates of sequestration are to be accurate. Unfortunately, there are no studies to date which have directly measured carbon sequestration of forests and wider non-turf areas on golf courses, making this a possible area of future research.


There is some disagreement within the turfgrass science community about whether golf courses should take credit for carbon sequestration in non-turfgrass areas of the course, given that these areas are often not directly managed by golf course managers and sometimes are not entirely owned by the golf course either. Because the turfgrass system is central to the operation of a golf course, many turf scientists advocate that the turf system - the emissions of managing the turf and the sequestration that the turf provides - should be the main focus for golf course operators.

The carbon balance of golf courses

The carbon balance of a golf course is equal to the weight of carbon emitted minus the weight of carbon sequestered (Figure 2). A carbon balance is most typically calculated over a calendar year but can also be calculated over longer time scales. For example, one could calculate the carbon balance of a golf course over its entire lifetime.

Carbon positive (+)
If more carbon is emitted than sequestered over a given time period, the golf course is said to be “ carbon positive ”.
Carbon negative (-)
If more carbon is sequestered than emitted, the golf course is said to be “ carbon negative ”.
Carbon neutral
If emissions and sequestration are approximately equal, the golf course is said to be “ carbon neutral ”. Net zero is another term that is synonymous with carbon neutrality.

Figure 2. The carbon balance of a golf facility



Study of golf course maintenance emissions in US and Europe (Part 2)

Figure 3 shows the cumulative carbon balance, emissions and sequestration on four US golf courses over a 150-year period. Cumulative carbon emissions were assumed to increase linearly over the golf courses' lifetime, meaning that the yearly emission rate stayed constant at approximately 160 Mg CO₂e yr⁻¹. Cumulative carbon sequestration was highest for the first 30 years

after a golf course is built, but then levels off almost completely after approximately 75 years.

The carbon balance of the golf courses in the study was negative for the first 30 years of operation, but then became positive after 30 years. The average age of the four golf courses in the study was 65 years, at which point the golf courses were approximately 5000 Mg CO₂e carbon positive. Unless emissions are greatly reduced, golf courses become increasingly carbon positive as they age.

For golf courses in the study to achieve carbon neutrality (ie net zero emissions) over an assumed 200-year life cycle, emission levels would need to be lowered from the current level of 160 Mg CO₂e yr⁻¹ to 30 Mg CO₂e yr⁻¹ - a reduction of over 5 times.





Figure 3. Average cumulative carbon emissions, sequestration and balance on four US golf courses.



Golf course carbon cycle.



Practical steps.

Transitioning to electric maintenance equipment and sourcing electricity from a low carbon source are the most effective ways for the majority of golf courses to reduce carbon emissions.

Minimising carbon emissions

Fortunately, the technology exists today to greatly reduce emission levels in golf course maintenance, which come primarily from fuel use for mowing and electricity use to power the maintenance building and irrigation pump. For golf courses that topdress fairways with sand, the emissions from mining, transporting and applying sand to fairways is likely the primary source of emissions at the golf course. However, for most golf courses the most effective way to reduce emissions is to reduce fuel use by transitioning to electric maintenance equipment, which will

reduce carbon emissions regardless of how the electricity is produced. However, the degree of emission reduction from electrification of maintenance equipment depends on the amount of carbon emitted per kWh of electricity generated.



Electrification of golf course maintenance equipment

Average emissions from electricity generation are displayed for 15 countries in Figure 4 below.


If a golf course is located in a country that emits more than 0.3kg CO₂e kWh⁻¹, it is advisable to consider sourcing electricity from a lower carbon source other than

the national electricity grid. This can be achieved by installing on-site solar panels, or by sourcing electricity from a low carbon source (ie hydroelectric, solar or wind farm). Many electricity companies offer such a service, which is often referred to as a green tariff programme.

A 2016 study found that 75-80% of mowing emissions are from mowing fairways and roughs (Tidåker et al., 2017). As a result, converting

to electric mowers in these areas of the course is most important. While there are not currently any traditional fairway and rough golf mowers on the market that are electric, there are GPS guided autonomous electric mowers available that can mow fairways and roughs. Golf courses are strongly encouraged to consider adopting this technology to reduce emissions from mowing.

Golf course managers only have a limited ability to increase the sequestration capacity of a golf course.



Maximising carbon sequestration

Unlike carbon emissions, over which golf course managers have a high degree of control, managers only have a limited ability to increase the sequestration capacity of a golf course. Soil disturbances, like aeration, lead to a reduction in soil organic matter levels and carbon emissions. Greens and tees are a relatively small area of the average golf course (usually less than 8-10%). Aerating these areas is commonly agronomically important and will not significantly affect the

golf course's carbon sequestration capacity. However, aerating fairways, which cover on average 30% of a golf course, will reduce a golf course's ability to sequester carbon. In addition, tilling and renovating existing turf surfaces will also lead to a loss of soil organic carbon and carbon emissions. Practices such as returning clippings and increasing the turfgrass cutting height (especially in roughs) will also lead to a slightly higher carbon sequestration capacity.

Reducing carbon emissions of golf course maintenance
1. Transition to electric machinery (eg mowers, utility vehicles etc)
2. Source electricity from low carbon sources (on-site renewable or green tariff)
3. Install on-site solar panels on the roof of maintenance buildings
4. Reduce volume of topdressing and areas topdressed if agronomically viable
5. Utilise N fertilisation in young turfgrass system (ie up to 10 years after planting) to develop a well established turfgrass system and then begin to reduce N rates as the turfgrass system ages. After 50 years, N rates should be greatly reduced in most circumstances.

Maximising carbon sequestration of golf course maintenance*
1. Maintain a healthy and actively growing perennial turfgrass system
2. Eliminate fairway aeration
3. Refrain from tilling and renovating existing turfgrass areas, especially those less than 50 years old
4. Return grass clippings to the turfgrass surface
5. Increase turfgrass cutting height where possible

*This is a list of generalised recommendations that should only be applied where agronomically viable and may not be appropriate for all golf courses. Carbon sequestration should be one of many factors considered when making agronomic decisions on a golf course.

Summary.

Anthropogenic carbon emissions have caused the Earth's global average surface temperature to rise by approximately 1°C since the Industrial Revolution. This rapid rise in temperatures presents myriad challenges, including but not limited to more extreme and unpredictable weather, droughts, flooding, food shortages, sea level rise and a decline in global biodiversity. All sectors of society, including the golf industry, must be engaged in mitigating the worst effects of climate change by reducing carbon emissions and increasing sequestration.

Carbon emissions in golf course maintenance primarily derive from burning fuels to power maintenance equipment and from the electricity used to power the maintenance facility and irrigation pump. Therefore, the most effective way for golf course managers to reduce emissions, is to electrify maintenance equipment and to source electricity from a low carbon source.

Golf course turfgrasses commonly sequester carbon from the atmosphere, but the rate of sequestration is heavily dependent on the historical land use and the age of the golf course. Some turfgrass species may sequester carbon more quickly than others, though this remains an active area of scientific research. While not the focus of this report, vegetation in non-turf areas of a golf course can also sequester carbon. However, the rate of sequestration in these areas is uncertain because of a lack of

research, making this a possible direction for future investigation.

If considering the carbon balance of golf course turfgrass areas over their lifetime, recent research suggests that most golf courses are carbon negative in the first 30 years of operation. However, after 30 years of operation, most golf courses become carbon positive and become increasingly carbon positive as they age.

This report outlines basic strategies for golf courses to reduce emissions and maximise sequestration. However, for golf courses to be carbon negative or carbon neutral over their lifetimes carbon emissions must be greatly reduced from current levels. Altering agronomic practices helps to increase carbon sequestration, but these changes alone will rarely make a significant impact on a golf courses carbon balance.



All sectors of society, including the golf industry, must be engaged in mitigating the worst effects of climate change by reducing carbon emissions and increasing sequestration.

References.

Bekken, M.A.H., & Soldat D.J. (2022). *Estimated Energy Use and Greenhouse Gas Emissions Associated with Golf Course Turfgrass Maintenance in the Northern USA.*

Golf Course Superintendents Association of America (GCSAA), (2017). *Golf Course Environmental Profile: Land Use Characteristics and Environmental Stewardship Programs on U.S. Golf Courses.* Environmental Institute for Golf.

Qian, Y., & Follet, R.F. (2002). *Assessing Soil Carbon Sequestration in Turfgrass Systems Using Long-Term Soil Testing Data.* Agronomy Journal, 94: 930-935.

Selhorst, A., & Lal, R. (2011). *Carbon budgeting in golf course soil of Central Ohio.* Urban Ecosyst. 14: 771-781.

Tidåker, P., Wesström, T., Kätterer, T. (2017). *Energy use and greenhouse gas emissions from turf management of two Swedish golf courses.* Urban Forestry & Urban Greening. 21: 80-87.





The R&A group of companies was formed in 2004 to take on The Royal and Ancient Golf Club of St Andrews' responsibilities for governing the Rules of Golf, staging The Open, golf's original championship, and developing the sport. The World Golf Museum in St Andrews is part of The R&A group.

Together The R&A and the USGA govern the sport of golf worldwide, operating in separate jurisdictions with a commitment to a single code for the Rules of Golf, Rules of Amateur Status and Equipment Standards. The R&A, through R&A Rules Ltd, governs the sport worldwide, outside of the United States and Mexico, on behalf of over 41 million golfers in 144 countries and with the consent of 159 organisations from amateur and professional golf.

The R&A has responsibility for running a series of world class amateur events and international matches in women's and girls' as well as men's and boys' golf. The R&A stages the AIG Women's Open and works with the DP World Tour to stage the Senior Open presented by Rolex.

The R&A is committed to investing £200 million over ten years in developing golf and supports the growth of the sport internationally, including the development and management of sustainable golf facilities. For more information, visit www.randa.org.



WORLD
GOLF
MUSEUM

RandA.org



**GOLF
COURSE
2030**